

LA-UR-14-27494

Approved for public release; distribution is unlimited.

Title: The Standard Model and Beyond with Ultracold Neutrons


Author(s): Broussard, Leah Jacklyn

Intended for: Invited colloquium at Case Western Reserve University

Issued: 2014-10-01 (rev.1)

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



The Standard Model and Beyond with Ultracold Neutrons

Leah Broussard

Los Alamos National Laboratory

September 25, 2014



Ultracold Neutrons

Class	Energy	Source
Fast	> 1 MeV	Fission reactions / Spallation
Slow	eV – keV	Moderation
Thermal	0.025 ev	Thermal equilibrium
Cold	$\mu\text{eV} - \text{meV}$	Cold moderation
Ultracold	$\leq 300 \text{ neV}$	Downscattering

How cold is Ultracold?

- Temperature < 4 mK
- Velocity < 8 m/s
- Usain Bolt ~ 12 m/s



Sensitive to Lab-Accessible Potentials

- Gravitational ($V = mgh$): 100 neV / meter
- Magnetic ($V = -\vec{\mu} \cdot \vec{B}$): 60 neV / Tesla

- Material $\left(V = \frac{2\pi\hbar^2 Nb}{m} \right) \begin{cases} {}^{58}\text{Ni} : & 335 \text{ neV} \\ \text{DLC} : & 250 \text{ neV} \\ \text{Cu} : & 170 \text{ neV} \end{cases}$





Ultracold Neutrons Live in Bottles

UCN video






LANSCE UCN Facility




UCN Source¹

- 800 MeV protons bombard a tungsten target to create spallation neutrons
- Neutrons are cooled in the polyethylene moderator and downscattered in a solid deuterium crystal to form UCN
- High densities observed: 50 UCN/cc at shield wall




¹Rev. Sci. Instrum. 84, 013304 (2013)

UCN Experimental Program




The Standard Model of Particle Physics

Standard Model is complete!



The Standard Model of Particle Physics

Standard Model is complete!




But...

- Why 3 generations?
- Why so many parameters?
- Why these masses?
- Why left-handed weak interaction?
- What is Dark Matter?
- Why more matter than anti-matter?
- Where is gravity?

The Standard Model of Particle Physics

Standard Model is complete!



But...

- Why 3 generations?
- Why so many parameters?
- Why these masses?
- Why left-handed weak interaction?
- What is Dark Matter?
- Why more matter than anti-matter?
- Where is gravity?


Are there more missing pieces?

- No more sure-thing theories!
- High Energy frontier (LHC) vs. Precision frontier (beta decay)
- High energy: Direct search for heavy particles
- Precision: Measure deviations from SM
- Complementary limits

The Weak Interaction

Described very well by the Standard Model

Example: β -decay



Quark Mixing (CKM)

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

- $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$

Nuclear β -decay exhibits influence from Strong Force

- Left-handed coupling: **Vector – AxialVector**
- $H = \frac{G_F V_{ud}}{\sqrt{2}} [\bar{e}(\gamma^\mu - \gamma^\mu \gamma_5) \nu_e \bar{u}(g_V \gamma_\mu + g_A \gamma_\mu \gamma_5) d]$
- Standard Model: $g_V = 1$, g_A free parameter

The Neutron Beta Decay Alphabet

Experimental Observables

- Angular correlations polarized decay¹:

$$\frac{dW}{dE_e d\Omega_e d\Omega_\nu} \propto p_e E_e (E_0 - E_e)^2 \left[1 + \mathbf{a} \frac{\vec{p}_e \cdot \vec{p}_\nu}{E_e E_\nu} + \mathbf{b} \frac{m_e}{E_e} + \langle \vec{\sigma}_n \rangle \cdot \left(\mathbf{A} \frac{\vec{p}_e}{E_e} + \mathbf{B} \frac{\vec{p}_\nu}{E_\nu} + \mathbf{D} \frac{\vec{p}_e \times \vec{p}_\nu}{E_e E_\nu} \right) \right]$$

- Lifetime:

$$\frac{1}{\tau_n} = W = K (G_F \mathbf{V}_{ud})^2 \left(1 + 3 \left(\frac{\mathbf{g_A}}{\mathbf{g_V}} \right)^2 \right) (1 + \Delta_R) f_n p_e E_e (E_0 - E_e)^2 \left[1 + m_e \mathbf{b} \frac{f_b}{f_n} \right]$$

- \mathbf{A} , \mathbf{a} + τ → \mathbf{V} , \mathbf{A} interactions
- \mathbf{B} , \mathbf{b} → \mathbf{S} , \mathbf{T} (BSM) interactions



Test CKM Unitarity: Extract \mathbf{V}_{ud}

- $\mathbf{a}_0 = \frac{1-\lambda^2}{1+3\lambda^2}$, $\mathbf{A}_0 = -2 \frac{\lambda(\lambda+1)}{1+3\lambda^2}$, $\mathbf{B}_0 = 2 \frac{\lambda(\lambda-1)}{1+3\lambda^2}$, $\tau = \frac{\text{constant}}{1+3\lambda^2}$
- \mathbf{A} Most sensitive to $\lambda = \frac{\mathbf{g_A}}{\mathbf{g_V}}$
- $\tau_n + \lambda \rightarrow$ extract CKM matrix element \mathbf{V}_{ud}




¹Phys. Rev. C **106** 517 (1957)


V_{ud} from Neutron Decay: A, a, and τ



- Superallowed Fermi $0^+ \rightarrow 0^+$ decays: V_{ud} at 0.02% level¹
- From neutron decay, require $\frac{\delta A}{A} \sim 0.1\% + \delta\tau \sim 0.35\text{s} \rightarrow V_{ud}$ at 0.02% level



¹ Ann. Phys. 525 443 (2013)





UCNA Collaboration

California Institute of Technology

R. Carr, B. W. Fillippone, M. P. Mendenhall, A. Perez-Galvan, R. Picker, S. Slutsky

Idaho State University

R. Rios, E. Tatar

Indiana University

A. T. Holley, C.-Y. Liu, D. Salvat

Institut Laue-Langevin

P. Geltenbort

Los Alamos National Laboratory

M. Blatnik, T. J. Bowles, L. Broussard, S. Clayton, S. Currie, G. Hogan, T. M. Ito, M. Makela, P. McGaughey, J. Mirabal, C. L. Morris, J. Ramsey, A. Saunders (co-spokesperson), S. Seestrom, E. Shaw, S. Sjue, W. Sondheim, T. Womack

Michigan State University

C. Wrede

North Carolina State University

E.B. Dees, J. Hoagland, R. Pattie, B. VornDick, A. Young, B. Zeck

Shanghai Jiao Tong University

J. L. Liu

Texas A&M University

D. Melconian

University of California, Los Angelos

K. Hickerson

University of Kentucky

M. Brown, S. Hasan, B. Plaster

University of Winnipeg

J. W. Martin


Virginia Tech

X. Ding, M. L. Pitt, R. B. Vogelaar






UCNA Experiment




$$W(E) \propto 1 + \frac{v}{c} \langle P \rangle A(E) \cos\theta$$

- 1 T field: $\langle \cos\theta \rangle = \pm \frac{1}{2}$
- P : limit depolarization





UCNA Experiment





- Reduce backscatter: low Z, field expansion
- Plastic Scintillator: β energy
- MWPC: β position info, suppress backgrounds, backscatter reconstruction
- Super-ratio: cancel loading/detector efficiencies

$$S(E) = \frac{N(E)_1^+ (N(E)_2^-)}{N(E)_1^- (N(E)_2^+)} \quad A(E) = \frac{1 - \sqrt{S(E)}}{1 + \sqrt{S(E)}}$$



UCNA Results: 2010 data set





- Mendenhall et al. *Phys Rev C* **87** 032501(R) (2013)
- 20M β -decay events
- $A_0 = 0.11972(55)_{\text{stat}}(98)_{\text{sys}}$



UCNA: 2011–2012 data sets and beyond

Improvement of UCNA Systematics (preliminary)

Uncertainty (%)	2010 dataset (Mendenhall 2013 PRC)	2011-2012 datasets (in analysis)	Next Step	Source of improvement
Statistics	+/- 0.46	+/- 0.40	+/- 0.28	Decay rate!
Depolarization	+0.67 +/- 0.56	+0.7 +/- 0.1	+0.7 +/- 0.1	Shutter+ ex situ
Backscatter	+1.36 +/- 0.34	+0.56 +/- 0.15	+0.56 +/- 0.15	Thin windows
Angle effect	-1.21 +/- 0.30	-0.8 +/- 0.2	-0.8 +/- 0.1	Windows+APD
Energy Reconstruction	+/- 0.31	+/- 0.08	+/- 0.08	Xenon + LED
Total Sys.	+/- 0.82	+/- 0.28	+/- 0.22	
Total	+/- 0.94	+/- 0.5	+/- 0.35	



UCN τ Collaboration

DePauw University

A. Komives

Hamilton College

G. Jones

Indiana University

E. R. Adamek, N. B. Callahan, W. Fox, C.-Y. Liu, D. J. Salvat, B. A. Slaughter, W. M. Snow, K. Solberg, J. Vanderwerp

JINR

E. I. Sharapov

Los Alamos National Laboratory

D. Barlow, M. Blatnik, L. J. Broussard, S. M. Clayton, T. M. Ito, M. Makela, J. Medina, D. J. Morley, C. L. Morris, R. W. Pattie, Jr., J. Ramsey, A. Saunders, S. J. Seestrom, S. K. L. Sjue, P. Walstrom, Z. Wang, T. L. Womack

North Carolina State University

C. Cude-Woods, E. B. Dees, B. Vorndick, A. R. Young, B. A. Zeck

University of California Los Angeles

K. P. Hickerson

Virginia Tech

X. Ding, B. Vogelaar

Tennessee Tech

A. T. Holley

The Neutron Lifetime τ

Beam of cold neutrons

- Count the dying

PHYSICAL REVIEW C 71, 055502 (2005)





FIG. 2. Experimental method for measuring lifetime by counting neutrons and trapped protons.

Bottle of ultracold neutrons

- Count the survivors



$$1/\tau_{\text{storage}} = 1/\tau_n + 1/\tau_{\text{loss}}$$

Challenge: neutron flux measurement

Beam vs. Bottle disagree by 8 s!



UCN τ Experiment

Material Bottles



- Extrapolate wall losses: large corrections!

Magneto-gravitational Trap


- World's largest permanent magnet array
- No material interactions!
- Asymmetric design: phase space mixing → no quasi-bound orbits




UCN τ Experiment



Ex-situ: fill and dump



Storage time > 10 hours!




UCN τ Experiment

Improve precision with *in-situ* measurement

- Capture UCNs: $^{51}V(n, \gamma)^{52}V$
- Detect: $^{52}V \rightarrow ^{52}Cr + \beta + \gamma$
- β scintillator + NaI γ detector assembly

UCN absorbed 10 \times faster than drained

V foil absorber height: important systematic check



The Weak Interaction Beyond V-A: B and b

What can cause a scalar/tensor contribution?



- SUSY, leptoquarks, ...
- Not yet observed at the LHC
- Need model independent extraction

Current limits on Scalars

- From Superallowed Fermi $0^+ \rightarrow 0^+$ decays¹
 $-1 \times 10^{-3} < C_S/C_V < 3 \times 10^{-3}$ (90% C.L.)

and Tensors

- From $\pi^+ \rightarrow e^+ \nu \gamma$ (PIBETA)²
 $-2 \times 10^{-3} < C_T/C_A < 2 \times 10^{-3}$ (90% C.L.)
- From combined neutron/nuclear beta decay data³
 $0 < C_T/C_A < 4 \times 10^{-3}$ (90% C.L.)



¹Phys. Rev. C **79** 055502 (2009)

²Phys. Rev. Lett. **103** 051802 (2009)

³arxiv:1306.2608v2 (2013)

BSM Scalar and Tensors in Neutron Decay

Access via b , B^1

$$\frac{d\Gamma}{dE_e d\Omega_e d\Omega_\nu} \propto w(E_e) \left(1 + \frac{m_e}{E_e} \bar{b} + \bar{a}(E_e) \frac{\vec{p}_e \cdot \vec{p}_\nu}{E_e E_\nu} + \bar{A}(E_e) \frac{\vec{\sigma}_n \cdot \vec{p}_e}{E_e} + \bar{B}(E_e) \frac{\vec{\sigma}_n \cdot \vec{p}_\nu}{E_\nu} + \dots \right)$$

- $\bar{B}(E_e) = \frac{2\bar{\lambda}(1+\bar{\lambda})}{1+3\bar{\lambda}^2} + \frac{m_e}{E_e} (b_\nu^{SM} + b_\nu^{BSM}) + c_0 + c_1 \frac{E_e}{M_N}$
- $\bar{\lambda} = \lambda(1 - 2\epsilon_R)$; $c_{0,1}$: recoil corrections
- $\bar{b} = b^{SM} + b^{BSM}$

New Physics in b^{BSM} , b_ν^{BSM}

$$\bullet b^{BSM} \sim 0.34g_S\epsilon_S - 5.22g_T\epsilon_T$$


$$b^{SM} = - \left(\frac{m_e}{M_N} \frac{1+2\mu_V+\lambda^2}{1+3\lambda^2} \right)$$

$$\bullet b_\nu^{BSM} \sim 0.44g_S\epsilon_S - 4.85g_T\epsilon_T$$

$$b_\nu^{SM} = - \left(\frac{m_e}{M_N} \frac{(1+\lambda)(\mu_V+\lambda)}{1+3\lambda^2} \right)$$

Experimental Determination of B

- Actually measure $B_{exp} = \frac{\bar{B}(E_e)}{1+b m_e/E_e}$
- $B_{exp} \propto b_\nu^{BSM} - b^{BSM}$





UCNB Collaboration

Los Alamos National Laboratory

L. Broussard, M. Makela, P. McGaughey, J. Mirabal, C. Morris, J. Ramsey, A. Saunders, S. Sjue, J. Wang, S. Wilburn, T. Womack

Caltech

C. Feng, B. Filippone

University of California, Los Angeles

K. Hickerson

Indiana University

A. T. Holley, D. J. Salvat, C.-Y. Liu

North Carolina State University

E.B. Dees, J. Hoagland, R. Pattie, B. VornDick, A. Young, B. Zeck

University of Kentucky

M. Brown, S. Hasan, B. Plaster

University of Virginia

A. Bacci, S. Baessler, D. Pocanic

Virginia Tech

D. Bravo, X. Ding, R. B. Vogelaar






Experimentally Determining B

Classic Approach


- Determine ν direction from e, p
- Count $N^{\beta p} = N^{\pm\pm}$ (β, p aligned vs. antialigned with σ_n)



$$B_{exp}(E) = \frac{N^{--}(E) - N^{++}(E)}{N^{--}(E) + N^{++}(E)}$$

$$C_{exp}(E) = \frac{(N^{++}(E) - N^{-+}(E)) - (N^{+-}(E) - N^{--}(E))}{(N^{++}(E) - N^{-+}(E)) + (N^{+-}(E) - N^{--}(E))}$$

- $\frac{\delta B}{B} = \frac{2.9}{\sqrt{N}}$: 0.1% in 1 month at 10Hz decay rate
- World average
 $B = 0.9807 \pm 0.0030(0.3\%)$





Isolating b_ν

Ratio of asymmetries¹

- $r \equiv \frac{\beta_e E_e}{E_\nu} \quad (r = 1 \text{ at } T_e = 236 \text{ keV})$

$$\alpha_p(E_e) = \frac{Q_{-\pm} - Q_{+\pm}}{Q_{-\pm} + Q_{+\pm}}$$

$$= \begin{cases} P \frac{\frac{2r}{3} A \beta_e + B \left(1 - \frac{r^2}{3}\right)}{2 \left(1 + b \frac{m_e}{E_e}\right)} & r < 1 \\ P \frac{A \beta_e \left(1 - \frac{1}{3r^2}\right) + \frac{2}{3r} B}{2 \left(1 + b \frac{m_e}{E_e}\right)} & r > 1 \end{cases}$$

$$\alpha_e(E_e) = \frac{Q_{\pm-} - Q_{\pm+}}{Q_{\pm-} + Q_{\pm+}}$$

$$= -\frac{1}{2 \left(1 + b \frac{m_e}{E_e}\right)} PA \beta_e$$


- Monte Carlo: $N = 10^8 \rightarrow$ probe b_ν at 3×10^{-3} level

Combined Fit

- $Q^{\pm\pm}: E_e, \alpha_e(E_e), \alpha_p(E_e)$
- Free parameters: λ, b, b_ν , normalization
- Use ϵ_S, ϵ_T dependence: $b = b_\nu + x, x = [-0.0005, 0.0003]$
- Monte Carlo: $N = 10^8 \rightarrow \lambda \sim 0.09\%, b \sim 2.2 \times 10^{-3}, b_\nu \sim 2.2 \times 10^{-3}$

¹B. Plaster, S. K. L. Sjue, A. R. Young, *in prep* (2014)

UCNB Experiment: Overview





UCNB Experiment

CHALLENGE: Detecting protons and betas in coincidence

Protons

- max $E < 800$ eV
- slow timing: $10 \mu\text{s}$ to 1 ms after decay
- Very low energy: detector deadlayer and noise important

Electrons

- max $E \approx 800$ keV
- fast timing: 10 ns
- problem: backscattering
 \rightarrow partial energy signal, direction


State of the art Si detectors



- 2 mm thick, 12 cm diameter active area
- Thin ($\sim 100 \text{ nm}$) deadlayer
- 128 pixels
- Characterization: Salas-Bacci *NIM A* (2014)



UCNB Experiment: Mount and Electrodes



- Re-entrant into spectrometer vacuum: detector located at 0.6 T flat region
- Detector-FET assembly in vacuum, rest of preamp in air
- Full DAQ biased: detector, preamps, digitizers
- Stable operation at 1 T and -30 kV for \sim 100 hours








UCNB: Current Status

Preamplifiers

- 8-ch: <15 keV noise demonstrated, ~ 3 keV energy resolution
- 24-ch: Now 20 ns rise time
- Successful demonstration with β -decay (Fall 2014) \rightarrow assemble 128 ch model





Neutron beta-decay proton-electron coincidences detected






UCNB partnership with Nab




- Planned for Spallation Neutron Source at Oak Ridge National Laboratory
- $$\frac{dW}{dE_e d\Omega_e d\Omega_\nu} \propto p_e E_e (E_0 - E_e)^2 \left[1 + \mathbf{a} \frac{\vec{p}_e \cdot \vec{p}_\nu}{E_e E_\nu} + \mathbf{b} \frac{m_e}{E_e} + \langle \vec{\sigma}_n \rangle \cdot \left(\mathbf{A} \frac{\vec{p}_e}{E_e} + \mathbf{B} \frac{\vec{p}_\nu}{E_\nu} + \mathbf{D} \frac{\vec{p}_e \times \vec{p}_\nu}{E_e E_\nu} \right) \right]$$
- Will measure **a** (similar sensitivity to λ as **A**) and **b** (zero in SM)
- No polarized neutron requirement
- Same detector/preamp technology

New effort to measure nEDM at LANL



Why EDM

- EDM nonzero \rightarrow T symmetry violation
- Matter-antimatter asymmetry? Source of CP violation?
- SM prediction: $d_n \sim 10^{-32} - 10^{-31}$ e-cm
- Current Limit: $d_n < 2.9 \times 10^{-26}$ e-cm (ILL)
- Constraints on EDM: best constraints on many BSM models






New effort to measure nEDM at LANL

Room temperature EDM

- Ramsey's separated oscillatory field method (similar to most precise result at ILL)
- EDM → frequency shift when \vec{E} -field reversed
- Goal: $\delta d_n \sim 10^{-27}$ e-cm
- ILL result: ~ 1 UCN/cc
- Requirement for improvement: 100 UCN/cc




LANL nEDM UCN Density Improvements




UCN Source

- Improved UCN source transition to guides
- Drive mechanism moved outside UCN volume
- Improvements to moderator cooling
- Geometry of source near W target



Proton Beam

- High current bursts every 30 s (presently every 5 s): Limit loss in SD₂



Transport to Experiment


- Current geometry suboptimal: Improve couplings, guide quality

Studies of Actinides

Typically 2 fragments emitted


- $A \sim 100$, $E \sim 100$ MeV, $\frac{v}{c} \sim 10\%$
- Range $\sim 10 \mu\text{m}$

^{235}U fission fragments



How does fission age material?

- Very energetic, heavy, charged particles
- How is energy deposited in material?
- Damage to the material?
- Near material surface: ejection of matter
- Differentiate between theoretical models



Why study nuclear material aging?

- Nuclear fuels
- Stockpile stewardship
- Lifetime of materials in space



Inducing Fission with Ultracold Neutrons

Experimental evidence

- Many previous measurements of sputtered atoms per fission
- **Significant disagreement in yield, distribution!**
- Key to differentiating models

New Technique for understanding sputtering

- Induce fission using Ultracold Neutrons
- Excellent control of neutron energy
- Very sensitive probe of fission as function of depth



LANL: Unique Position for work


- LANSCE: one of world's brightest sources of UCN
- Expertise in fabrication and analysis of actinide targets



Predictions for UCN-induced Fission in Uranium

Uncharted energy regime

- 3 orders of magnitude lower than ever explored
- Very high theoretical cross section
 $\sigma \sim \frac{1}{v}$
- 300 neV UCN: 2.16×10^5 barn



Control depth of fission event

Range of UCN in uranium (μm)						
	DU	NatU	SEU	LEU	HEU	VHEU
% ²³⁵ U	0.2	0.7	2	5	20	100
200 neV	118	66	312	13	4	0.8
300 neV	144	81	38	17	4.5	0.9
400 neV	191	101	45	20	5	1

First Observations of Fission from UCN

Detection chamber



- Cylindrical ion chamber

^{238}U

- 2.25 cm diameter, 1 mm thick disk of DU ($\sim 0.2\% \ ^{235}\text{U}$)
- Rate: $(1.3 \pm 0.8) \times 10^{-4}$ fission/UCN

^{235}U


- 2.2 cm diameter, 1 mm thick disk of HEU ($> 80\% \ ^{235}\text{U}$)
- Rate: $(1.90 \pm 0.02) \times 10^{-2}$ fission/UCN



UCN-energy dependence of fission rates


DU sample

- Electropolished, 2.25 cm diameter, 1 mm thick disk
- Measured fission rate decreases with UCN energy



^{235}U sample


- Thin sample: 30 μg heavily oxidized ^{235}U on tape
- Measured fission rate increases as UCN energy increases



Sputtering from UCN-induced fission


Exposure to Silicon wafer

- 1" diameter, 475 μm thick, polished wafer
- Exposed to 2.2 cm diameter, 1 mm thick electropolished DU disk
- 3×10^7 total UCN in chamber
- 0.18 μg ^{238}U collected on wafer
- Analyze with scanning probe microscopy



Exposure to Ni cylindrical foil


- 0.005" Ni foil, 1.15" diameter, 2.835" height
- UCN bottle: Ni material potential ~ 300 neV
- Exposed to DU disk and 30 μg ^{235}U thin sample



Sputtering on Nickel foil


Expose to DU disk and HEU film

- Each sample exposed to $\sim 10^8$ UCN
- Determine yield from α decay rate in ion chamber
- Can distinguish α 's from ^{238}U , ^{235}U decay



Angular distribution

- Mask vertical sections of foil
- Results from DU foil
- Distribution \sim isotropic





UCN Experimental Program

UCNA

- Expect to announce unblinded result end of this year

UCNB

- This fall: Detector characterization with ~35 pixels
- Next year: Full instrumentation of 128 pixels

UCN τ

- Moving to a 1 second accuracy measurement possibly this year

nEDM

- This fall: construct and test HV prototype chamber
- Next year: Install and test new source

Actinide Studies

- Install new beamline
- Continue detailed fission and sputtering characterizations



Opportunities for Students at LANL

Student Programs

- Undergraduate summer internships
- Post Baccalaureate programs (one year appointment)
- Post master's program (one year appointment)
- Graduate Research Assistants

Postdoctoral Program

- Postdoctoral Research Associates (2+1 year appointments)
- Director's, Agnew, Metropolis, Intelligence-Community Fellowships
- Distinguished Postdoctoral Fellowships (Oppenheimer, Feynman, Reines, Curie)

Seaborg Institute

- Research in nuclear science, especially actinides
- Nuclear Forensics Undergraduate Summer School
- Seaborg Student Research Fellowships
- Seaborg Postdoctoral Fellowship

